

UCB, Inc. Is Committed to Making FINTEPLA Affordable and Accessible

Patients get extra support when they need it. Once you and your patient's caregiver have decided that FINTEPLA is right for their loved one, they may have questions about financial assistance programs. UCB is committed to making FINTEPLA affordable and accessible for patients through financial assistance programs, including the ONWARD Copay Assistance Program, the ONWARD Echocardiogram Copay Support Program, and the ONWARD Support Program. Guide them to UCB, Inc. at 1-844-599-2273, where they can be matched with a single, dedicated Care Coordinator, who will help them navigate their financial options.

For families who are insured by their employer or self-insured and need help with their FINTEPLA out-of-pocket costs

ONWARD Copay Assistance Program

Through this program, families can pay as little as \$0 copays* for FINTEPLA and associated echocardiograms. The UCB commitment to patients is that they won't have to pay any more than \$25 in out-of-pocket copays for FINTEPLA. See eligibility requirements below.



Pay as little as **\$0** in out-of-pocket costs for **FINTEPLA** and **echocardiograms**.

Most families will not pay more than \$25 in out-of-pocket copays for FINTEPLA*

Families should understand all of their options for funding treatment. Whether they are insured by their employer, self-insured, covered by Medicaid or other government insurance, or not insured at all, a Care Coordinator will work with caregivers to understand their cost of treatment and connect them with financial assistance programs.

***ONWARD Copay Assistance Program Eligibility Requirements:**

Patients must have a FINTEPLA prescription from a licensed prescriber and be a resident of the United States or the Commonwealth of Puerto Rico, with a valid mailing address (no PO boxes). The Copay Assistance Program is not health insurance and is not valid for prescriptions that are eligible to be reimbursed, in whole or in part, by Medicaid, Medicare, or other government healthcare programs. Certain state restrictions may apply. UCB reserves the right to change or discontinue this offer without notice. Full terms and conditions can be found at [Fintepla.com/savings-terms](https://fintepla.com/savings-terms).



Please see Important Safety Information on next page and full [Prescribing Information](#), including Boxed Warning, for additional Important Safety Information on FINTEPLA.

INDICATIONS AND USAGE

FINTEPLA is indicated for the treatment of seizures associated with Dravet syndrome (DS) and Lennox-Gastaut syndrome (LGS) in patients 2 years of age and older.

IMPORTANT SAFETY INFORMATION

BOXED WARNING: VALVULAR HEART DISEASE and PULMONARY ARTERIAL HYPERTENSION

- **There is an association between serotonergic drugs with 5-HT_{2B} receptor agonist activity, including fenfluramine (the active ingredient in FINTEPLA), and valvular heart disease and pulmonary arterial hypertension.**
- **Echocardiogram assessments are required before, during, and after treatment with FINTEPLA.**
- **FINTEPLA is available only through a restricted program called the FINTEPLA REMS.**

CONTRAINDICATIONS

FINTEPLA is contraindicated in patients with hypersensitivity to fenfluramine or any of the excipients in FINTEPLA and with concomitant use, or within 14 days of the administration, of monoamine oxidase inhibitors because of an increased risk of serotonin syndrome.

WARNINGS AND PRECAUTIONS

Valvular Heart Disease and Pulmonary Arterial Hypertension (see Boxed Warning):

Because of the association between serotonergic drugs with 5-HT_{2B} receptor agonist activity, including fenfluramine (the active ingredient in FINTEPLA), and valvular heart disease (VHD) and pulmonary arterial hypertension (PAH), cardiac monitoring via echocardiogram is required prior to starting treatment, during treatment, and after treatment with FINTEPLA concludes. Cardiac monitoring via echocardiogram can aid in early detection of these conditions. In clinical trials for DS and LGS of up to 3 years in duration, no patient receiving FINTEPLA developed VHD or PAH.

Monitoring: Prior to starting treatment, patients must undergo an echocardiogram to evaluate for VHD and PAH. Echocardiograms should be repeated every 6 months, and once at 3-6 months post treatment with FINTEPLA.

The prescriber must consider the benefits versus the risks of initiating or continuing treatment with FINTEPLA if any of the following signs are observed via echocardiogram: valvular abnormality or new abnormality; VHD indicated by mild or greater aortic regurgitation or moderate or greater mitral regurgitation, with additional characteristics of VHD (eg, valve thickening or restrictive valve motion); PAH indicated by elevated right heart/pulmonary artery pressure (PASP >35 mm Hg).

FINTEPLA REMS Program (see Boxed Warning): FINTEPLA is available only through a restricted distribution program called the FINTEPLA Risk Evaluation and Mitigation Strategy

Warnings and Precautions (cont.)

(REMS) Program. Prescribers must be certified by enrolling in the FINTEPLA REMS. Prescribers must counsel patients receiving FINTEPLA about the risk of VHD and PAH, how to recognize signs and symptoms of VHD and PAH, the need for baseline (pretreatment) and periodic cardiac monitoring via echocardiogram during FINTEPLA treatment, and cardiac monitoring after FINTEPLA treatment. Patients must enroll in the FINTEPLA REMS and comply with ongoing monitoring requirements. The pharmacy must be certified by enrolling in the FINTEPLA REMS and must only dispense to patients who are authorized to receive FINTEPLA. Wholesalers and distributors must only distribute to certified pharmacies. Further information is available at www.FinteplaREMS.com or by telephone at 1-877-964-3649.

Decreased Appetite and Decreased Weight: FINTEPLA can cause decreases in appetite and weight. Decreases in weight appear to be dose related. Approximately half of the patients with LGS and most patients with DS resumed the expected measured increases in weight during the open-label extension studies. Weight should be monitored regularly during treatment with FINTEPLA, and dose modifications should be considered if a decrease in weight is observed.

Somnolence, Sedation, and Lethargy: FINTEPLA can cause somnolence, sedation, and lethargy. Other central nervous system (CNS) depressants, including alcohol, could potentiate these effects of FINTEPLA. Prescribers should monitor patients for somnolence and sedation and should advise patients not to drive or operate machinery until they have gained sufficient experience on FINTEPLA to gauge whether it adversely affects their ability to drive or operate machinery.

Suicidal Behavior and Ideation: Antiepileptic drugs (AEDs), including FINTEPLA, increase the risk of suicidal thoughts or behaviors in patients taking these drugs for any indication. Patients treated with an AED for any indication should be monitored for the emergence or worsening of depression, suicidal thoughts or behaviors, or any unusual changes in mood or behavior.

Anyone considering prescribing FINTEPLA or any other AED must balance the risk of suicidal thoughts or behaviors with the risks of untreated illness. Epilepsy and many other illnesses for which AEDs are prescribed are themselves associated with morbidity and mortality and an increased risk of suicidal thoughts and behaviors. Should suicidal thoughts and behaviors emerge during treatment, consider whether the emergence of these symptoms in any given patient may be related to the illness being treated.

Withdrawal of Antiepileptic Drugs: As with most AEDs, FINTEPLA should generally be withdrawn gradually because of the risk of increased seizure frequency and status epilepticus.

Continued on next page.

IMPORTANT SAFETY INFORMATION (cont.)

Warnings and Precautions (cont.)

If withdrawal is needed because of a serious adverse reaction, rapid discontinuation can be considered.

Serotonin Syndrome: Serotonin syndrome, a potentially life-threatening condition, may occur with FINTEPLA, particularly during concomitant administration of FINTEPLA with other serotonergic drugs, including, but not limited to, selective serotonin-norepinephrine reuptake inhibitors (SNRIs), selective serotonin reuptake inhibitors (SSRIs), tricyclic antidepressants (TCAs), bupropion, triptans, dietary supplements (eg, St. John's Wort, tryptophan), drugs that impair metabolism of serotonin (including monoamine oxidase inhibitors [MAOIs], which are contraindicated with FINTEPLA), dextromethorphan, lithium, tramadol, and antipsychotics with serotonergic agonist activity. Patients should be monitored for the emergence of signs and symptoms of serotonin syndrome, which include mental status changes (eg, agitation, hallucinations, coma), autonomic instability (eg, tachycardia, labile blood pressure, hyperthermia), neuromuscular signs (eg, hyperreflexia, incoordination), and/or gastrointestinal symptoms (eg, nausea, vomiting, diarrhea). If serotonin syndrome is suspected, treatment with FINTEPLA should be stopped immediately and symptomatic treatment should be started.

Increase in Blood Pressure: FINTEPLA can cause an increase in blood pressure. Rare cases of significant elevation in blood pressure, including hypertensive crisis, has been reported in adult patients treated with fenfluramine, including patients without a history of hypertension. In clinical trials for DS and LGS of up to 3 years in duration, no pediatric or adult patient receiving FINTEPLA developed hypertensive crisis. Monitor blood pressure in patients treated with FINTEPLA.

Glaucoma: Fenfluramine can cause mydriasis and can precipitate angle closure glaucoma. Consider discontinuing treatment with FINTEPLA in patients with acute decreases in visual acuity or ocular pain.

ADVERSE REACTIONS

The most common adverse reactions observed in DS studies (incidence at least 10% and greater than placebo) were decreased appetite; somnolence, sedation, lethargy; diarrhea; constipation; abnormal echocardiogram; fatigue, malaise, asthenia; ataxia, balance disorder, gait disturbance; blood pressure increased; drooling, salivary hypersecretion; pyrexia; upper respiratory tract infection; vomiting; decreased weight; fall; status epilepticus.

Adverse Reactions (cont.)

The most common adverse reactions observed in the LGS study (incidence at least 10% and greater than placebo) were diarrhea; decreased appetite; fatigue; somnolence; vomiting.

DRUG INTERACTIONS

Strong CYP1A2, CYP2B6, or CYP3A Inducers: Coadministration with strong CYP1A2, CYP2B6, or CYP3A inducers will decrease fenfluramine plasma concentrations. If coadministration of a strong CYP1A2, CYP2B6, or CYP3A inducer with FINTEPLA is necessary, monitor the patient for reduced efficacy and consider increasing the dosage of FINTEPLA as needed. If a strong CYP1A2, CYP2B6, or CYP3A inducer is discontinued during maintenance treatment with FINTEPLA, consider gradual reduction in the FINTEPLA dosage to the dose administered prior to initiating the inducer.

Strong CYP1A2 or CYP2D6 Inhibitors: Coadministration with strong CYP1A2 or CYP2D6 inhibitors will increase fenfluramine plasma concentrations. If FINTEPLA is coadministered with strong CYP1A2 or CYP2D6 inhibitors, the maximum daily dosage of FINTEPLA is 20 mg. If a strong CYP1A2 or CYP2D6 inhibitor is discontinued during maintenance treatment with FINTEPLA, consider gradual increase in the FINTEPLA dosage to the dose recommended without CYP1A2 or CYP2D6 inhibitors. If FINTEPLA is coadministered with stiripentol and a strong CYP1A2 or CYP2D6 inhibitor, the maximum daily dosage of FINTEPLA is 17 mg.

USE IN SPECIFIC POPULATIONS

In patients with severe impairment of kidney function (estimated glomerular filtration rate [eGFR]) 15 to 29 mL/min/1.73m², dosage adjustments are recommended. FINTEPLA has not been studied in patients with kidney failure (eGFR <15 mL/min/1.73m²).

Combined molar exposures of fenfluramine and norfenfluramine were increased in subjects with various degrees of hepatic impairment (Child-Pugh Class A, B, and C), necessitating a dosage adjustment in these patients.

To report SUSPECTED ADVERSE REACTIONS, contact UCB, Inc. at 1-844-599-2273 or FDA at 1-800-FDA-1088 or www.fda.gov/medwatch.

Please see full [Prescribing Information](#), including **Boxed Warning**, for additional Important Safety Information on FINTEPLA.